

Supplementary Online Content

Vega Potler NJ, Zhang J, Hackley B, et al. Persistence of emotional distress in unaccompanied migrant children and adolescents primarily from the Northern Triangle of Central America. *JAMA Netw Open*. 2023;6(6):e2318977. doi:10.1001/jamanetworkopen.2023.18977

eTable. Comparison of Unaccompanied Migrant Children Who Completed Only Initial vs Initial and Follow-up RHS-15

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Comparison of Unaccompanied Migrant Children Who Completed Only Initial vs Initial and Follow-up RHS-15

Variable	Number of RHS-15		P-value
	Completed only Initial RHS-15 (n = 108)	Completed Follow-up RHS-15 (n = 68)	
Age at initial RHS-15 (Years), mean (SD)	17.2 (2.2)	16.4 (1.7)	.01
Sex, n (%)			.02
Male	84 (77.8)	42 (61.8)	
Female	24 (22.2)	26 (38.2)	
Time in US at initial RHS-15, n (%)			.01
1 year or less	28 (33.3)	35 (57.4)	
>1 to 2 years	26 (31.0)	16 (26.2)	
Over 2 years	30 (35.7)	10 (16.4)	
Initial RHS-15 above the cutoff, n (%)			.76
Negative (below)	47 (43.5)	28 (41.2)	
Positive (above)	61 (56.5)	40 (58.8)	

Abbreviation: RHS-15, 15-item Refugee Health Screener